

APPLICATION

FOR

UNITED STATES LETTERS PATENT

BY

CARSTEN SJOEHOLM

AND

PETER RAHBK OESTERGAARD

FOR

USE OF ACID-STABLE PROTEASES IN ANIMAL FEED

Use of Acid-stable Proteases in Animal Feed.

Technical Field

The present invention relates to the use of acid-stable
5 proteases in animal feed (*in vivo*), and to the use of such
proteases for treating vegetable proteins (*in vitro*).

Proteins are essential nutritional factors for animals
and humans. Most livestock and many human beings get the
necessary proteins from vegetable protein sources. Important
10 vegetable protein sources are e.g. oilseed crops, legumes and
cereals.

When e.g. soybean meal is included in the feed of mono-
gastric animals such as pigs and poultry, a significant
proportion of the soybean meal solids is not digested. E.g.,
15 the apparent ileal protein digestibility in piglets and
growing pigs is only around 80%.

The stomach of mono-gastric animals and many fish
exhibits a strongly acidic pH. Most of the protein digestion,
however, occurs in the small intestine. A need therefore
20 exists for an acid-stable protease that can survive passage of
the stomach.

Background Art

The use of proteases in animal feed, or to treat
25 vegetable proteins, is known from the following documents:

WO95/28850 discloses i.a. an animal feed additive
comprising a phytase and a proteolytic enzyme. Various
proteolytic enzymes are specified at p. 7.

WO96/05739 discloses an enzyme feed additive comprising
30 xylanase and a protease. Suitable proteases are listed at p.
25.

US 3966971 discloses a process of obtaining protein from
vegetable protein sources by treatment with an acid-stable protease.

and optionally a proteolytic enzyme. Suitable proteases are specified in column 2.

US 4073884, US 5047240, US 3868448, US 3823072, and US 3683069 describe protease preparations derived from various strains of Streptomyces and their use in animal feed.

These proteases, however, are not acid-stable and/or are not homologous to the proteases described herein.

Brief Description of the Invention

10 Proteases have now been identified which are found to be very acid-stable, and expectedly of an improved performance in animal feed.

Brief Description of Drawings

15 The present invention is further illustrated by reference to the accompanying drawings, in which:

Fig. 1 shows pH-stability curves, viz. residual protease activity of five proteases (two acid-stable proteases derived from Nocardiosis, and three reference proteases (Sub.Novo, 20 and Sub.Novo(Y217L), both derived from Bacillus amylolique-faciens, and SAVINASE™) after incubation for 2 hours, at a temperature of 37°C, and at pH-values in the range of pH 2 to pH 11; the activity is relative to residual activity after a 2 hour incubation at pH 9.0, and 5°C;

25 Fig. 2 shows pH-activity curves, viz. protease activity between pH 3 and pH 11, relative to the protease activity at pH-optimum, of the same five proteases; and

Fig. 3 shows temperature-activity curves at pH 9.0, viz. protease activity at pH 9.0 between 15°C and 80°C, relative to 30 protease activity at the optimum temperature, of the same five proteases.

4. A hydrolase peptide bond has protease activity. Proteases are also called e.g. peptidases, proteinases, peptide

Preferred proteases for use according to the invention are of the endo-type that act internally in polypeptide chains (endopeptidases). Endopeptidases show activity on N- and C-terminally blocked peptide substrates that are relevant for the specificity of the protease in question.

Included in the above definition of protease are any enzymes belonging to the EC 3.4 enzyme group (including each of the thirteen sub-subclasses thereof) of the EC list (Enzyme Nomenclature 1992 from NC-IUBMB, 1992), as regularly supplemented and updated, see e.g. the World Wide Web (WWW) at <http://www.chem.qmw.ac.uk/iubmb/enzyme/index.html>.

Proteases are classified on the basis of their catalytic mechanism into the following groupings: serine proteases (S), cysteine proteases (C), aspartic proteases (A), metalloproteases (M), and unknown, or as yet unclassified, proteases (U), see Handbook of Proteolytic Enzymes, A.J.Barrett, N.D.Rawlings, J.F.Woessner (eds), Academic Press (1998), in particular the general introduction part.

The Nocardiosis proteases disclosed herein are serine proteases.

In a particular embodiment the proteases for use according to the invention are serine proteases. The term serine protease refers to serine peptidases and their clans as defined in the above Handbook. In the 1998 version of this handbook, serine peptidases and their clans are dealt with in chapters 1-175. Serine proteases may be defined as peptidases in which the catalytic mechanism depends upon the hydroxyl group of a serine residue acting as the nucleophile that attacks the peptide bond. Examples of serine proteases for use according to the invention are proteases of Clan SA, e.g. Family S2 (Streptogrisin), e.g. Sub-family S2A (alpha-lytic protease), as defined in the above Handbook.

Assay pH and assay temperature are likewise to be adapted to

5, 6, 7, 8, 9, 10, or 11. Examples of assay-temperatures are 25, 30, 35, 37, 40, 45, 50, 55, 60, 65, or 70°C.

Examples of protease substrates are casein, and pNA-substrates, such as Suc-AAPF-pNA (available e.g. from Sigma S-5 7388). The capital letters in this pNA-substrate refers to the one-letter amino acid code. Another example is Protazyme AK (azurine-dyed crosslinked casein prepared as tablets by Megazyme T-PRAK). For pH-activity and pH-stability studies, the pNA-substrate is preferred, whereas for temperature-10 activity studies, the Protazyme AK substrate is preferred.

Examples of protease assays are described in the experimental part.

There are no limitations on the origin of the protease for use according to the invention. Thus, the term protease 15 includes not only natural or wild-type proteases, but also any mutants, variants, fragments etc. thereof exhibiting protease activity, as well as synthetic proteases, such as shuffled proteases, and consensus proteases. Such genetically engineered proteases can be prepared as is generally known in 20 the art, e.g. by Site-directed Mutagenesis, by PCR (using a PCR fragment containing the desired mutation as one of the primers in the PCR reactions), or by Random Mutagenesis. The preparation of consensus proteins is described in e.g. EP 897985.

25 Examples of acid-stable proteases for use according to the invention are

(i) the proteases derived from *Nocardiopsis* sp. NRRL 18262, and *Nocardiopsis alba*;

(ii) proteases of at least 60, 65, 70, 75, 80, 85, 90, 30 or at least 95% amino acid identity to any of the proteases of (i);

(iii) proteases of at least 60, 65, 70, 75, 80, 85, 90, or at least 95% amino acid identity to any of the proteases of (i) or (ii);

For determining percentage identity, any computer program known in the art can be used. Examples of such

10

D.

20

30

Examples of microorganisms are bacteria and fungi.

Order: *Ascomycetes*, Class: *Fungi*, Subclass: *Xylariomycetes*, Order: *Xylariales*, Family: *Xylariaceae*, Genus: *Xylaria*, Species: *Xylaria* sp. 1.

exhibiting protease activity. This taxonomy is on the basis of Bergey's Manual of Systematic Bacteriology, 2nd edition, 2000, Springer (preprint: Road Map to Bergey's).

Further examples of microorganisms are fungi, such as yeast or filamentous fungi.

In another embodiment the protease is a plant protease. An example of a protease of plant origin is the protease from the sarcocarp of melon fruit (Kaneda et al, J.Biochem. 78, 1287-1296 (1975)).

The term animal includes all animals, including human beings. Examples of animals are non-ruminants, and ruminants, such as cows, sheep and horses. In a particular embodiment, the animal is a non-ruminant animal. Non-ruminant animals include mono-gastric animals, e.g. pigs or swine (including, but not limited to, piglets, growing pigs, and sows); poultry such as turkeys and chicken (including but not limited to broiler chicks, layers); young calves; and fish (including but not limited to salmon).

The term feed or feed composition means any compound, preparation, mixture, or composition suitable for, or intended for intake by an animal.

In the use according to the invention the protease can be fed to the animal before, after, or simultaneously with the diet. The latter is preferred.

In the present context, the term acid-stable means, that the protease activity of the pure protease enzyme, in a dilution corresponding to $A_{280} = 1.0$, and following incubation for 2 hours at 37°C in the following buffer:

100mM succinic acid, 100mM HEPES, 100mM CHES,
100mM CABS, 1mM CaCl_2 , 150mM KCl, 0.01% Triton® X-100, pH 3.5,

is at least 40% of the reference activity as measured using

In particular embodiments of the above acid stability definition, the protease activity is at least 45, 50, 55, 60,

The term reference activity refers to the protease activity of the same protease, following incubation in pure form, in a dilution corresponding to $A_{280} = 1.0$, for 2 hours at 5°C in the following buffer: 100mM succinic acid, 100mM HEPES, 5 100mM CHES, 100mM CABS, 1mM CaCl_2 , 150mM KCl, 0.01% Triton® X-100, pH 9.0, wherein the activity is determined as described above.

In other words, the method of determining acid-stability comprises the following steps:

- 10 a) The protease sample to be tested (in pure form, $A_{280} = 1.0$) is divided in two aliquots (I and II);
- b) Aliquot I is incubated for 2 hours at 37°C and pH 3.5;
- c) Residual activity of aliquot I is measured (pH 9.0 15 and 25°C);
- d) Aliquot II is incubated for 2 hours at 5°C and pH 9.0;
- e) Residual activity of aliquot II is measured (pH 9.0 and 25°C);
- 20 f) Percentage residual activity of aliquot I relative to residual activity of aliquot II is calculated.

Alternatively, in the above definition of acid-stability, the step b) buffer pH-value may be 1.0, 1.5, 2.0, 2.5, 3.0, 3.1, 3.2, 3.3, or 3.4.

- 25 In other alternative embodiments of the above acid-stability definition relating to the above alternative step b) buffer pH-values, the residual protease activity as compared to the reference, is at least 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, or at least 97%.

- 30 In alternative embodiments, pH values of 6.0, 6.5, 7.0, 7.5, 8.0, or 8.5 can be applied for the step d) buffer.

In the above acid-stability definition, the term A

refers to the activity of the protease in the reference buffer.

And in the above acid stability definition, the term

refers to the activity of the protease in the test buffer.

Coomassie-stained SDS-PAGE gel is measured to have at least 95% of its scan intensity in the band corresponding to said protease (see Example 2A). In the alternative, the A_{280}/A_{260} ratio is above or equal to 1.50, 1.60, 1.65, 1.70, 1.75, 1.80, 5 1.85, or above or equal to 1.90.

However, for the uses according to the invention, the protease need not be that pure; it may e.g. include other enzymes, even other proteases, in which case it could be termed a protease preparation. Nevertheless, a well-defined 10 protease preparation is advantageous. For instance, it is much easier to dose correctly to the feed a protease that is essentially free from interfering or contaminating other proteases. The term dose correctly refers in particular to the objective of obtaining consistent and constant results, and 15 the capability of optimising dosage based upon the desired effect.

In a particular embodiment, the protease, in the form in which it is added to the feed, or when being included in a feed additive, is well-defined. Well-defined means that the 20 protease preparation is at least 50% pure as determined by Size-exclusion chromatography (see Example 8).

In other particular embodiments the protease preparation is at least 60, 70, 80, 85, 88, 90, 92, 94, or at least 95% pure as determined by this method.

25 In the alternative, the term well-defined means, that a fractionation of the protease preparation on an appropriate Size-exclusion column reveals only one major protease component.

The skilled worker will know how to select an 30 appropriate Size-exclusion chromatography column. He might start by fractionating the preparation on e.g. a HiLoad26/60 Superdex75pg column from Amersham Pharmacia Biotech (see

particular example 8). The skilled worker will know how to select the sample volume. By simple and common trial and-error methods he will find the appropriate sample volume for the column and the preparation.

(clear separation of peaks), on the basis of which the purity calculation is performed as described in Example 8.

The protease preparation can be (a) added directly to the feed (or used directly in the treatment process of vegetable proteins), or (b) it can be used in the production of one or more intermediate compositions such as feed additives or premixes that is subsequently added to the feed (or used in a treatment process). The degree of purity described above refers to the purity of the original protease preparation, whether used according to (a) or (b) above.

Protease preparations with purities of this order of magnitude are in particular obtainable using recombinant methods of production, whereas they are not so easily obtained and also subject to a much higher batch-to-batch variation when the protease is produced by traditional fermentation methods.

Such protease preparation may of course be mixed with other enzymes.

In one particular embodiment, the protease for use according to the invention, besides being acid-stable, also has a pH-activity optimum close to neutral.

The term pH-activity optimum close to neutral means one or more of the following: That the pH-optimum is in the interval of pH 6.0-11.0, or pH 7.0-11.0, or pH 6.0-10.0, or pH 7.0-10.0, or pH 8.0-11.0, or pH 8.0-10.0 (see Example 2B and Fig. 2 herein).

In another particular embodiment, the protease for use according to the invention, besides being acid-stable, is also thermostable.

The term thermostable means one or more of the following: That the temperature optimum is at least 50°C, 52°C,

According to the invention is capable of stabilizing vegetable proteins according to the *in vitro* model of Example

The term vegetable proteins as used herein refers to any compound, composition, preparation or mixture that includes at least one protein derived from or originating from a vegetable, including modified proteins and protein-
5 derivatives. In particular embodiments, the protein content of the vegetable proteins is at least 10, 20, 30, 40, 50, or 60% (w/w).

Vegetable proteins may be derived from vegetable protein sources, such as legumes and cereals, for example materials
10 from plants of the families Fabaceae (Leguminosae), Cruciferaceae, Chenopodiaceae, and Poaceae, such as soy bean meal, lupin meal and rapeseed meal.

In a particular embodiment, the vegetable protein source is material from one or more plants of the family Fabaceae,
15 e.g. soybean, lupine, pea, or bean.

In another particular embodiment, the vegetable protein source is material from one or more plants of the family Chenopodiaceae, e.g. beet, sugar beet, spinach or quinoa.

Other examples of vegetable protein sources are
20 rapeseed, and cabbage.

Soybean is a preferred vegetable protein source.

Other examples of vegetable protein sources are cereals such as barley, wheat, rye, oat, maize (corn), rice, and sorghum.

25 The treatment according to the invention of vegetable proteins with at least one acid-stable protease results in an increased solubilisation of vegetable proteins.

The following are examples of % solubilised protein obtainable using the proteases of the invention: At least
30 74.0%, 74.5%, 75.0%, 75.5%, 76.0%, 76.5%, 77.0%, or at least 77.5%, reference being had to the *in vitro* model of Example 4 herein.

The protease-mediated release of protein from other components of the usually complex natural compositions such as

amount of soluble proteins, by reference to a sample with no protease treatment (see Example 4 herein).

In a particular embodiment of a treatment process the protease(s) in question is affecting (or acting on, or exerting its solubilising influence on the vegetable proteins or protein sources. To achieve this, the vegetable protein or protein source is typically suspended in a solvent, eg an aqueous solvent such as water, and the pH and temperature values are adjusted paying due regard to the characteristics of the enzyme in question. For example, the treatment may take place at a pH-value at which the relative activity of the actual protease is at least 50, or 60, or 70, or 80 or 90%. Likewise, for example, the treatment may take place at a temperature at which the relative activity of the actual protease is at least 50, or 60, or 70, or 80 or 90% (these relative activities being defined as in Example 2 herein). The enzymatic reaction is continued until the desired result is achieved, following which it may or may not be stopped by inactivating the enzyme, e.g. by a heat-treatment step.

In another particular embodiment of a treatment process of the invention, the protease action is sustained, meaning e.g. that the protease is added to the vegetable proteins or protein sources, but its solubilising influence is so to speak not switched on until later when desired, once suitable solubilising conditions are established, or once any enzyme inhibitors are inactivated, or whatever other means could have been applied to postpone the action of the enzyme.

In one embodiment the treatment is a pre-treatment of animal feed or vegetable proteins for use in animal feed, i.e. the proteins are solubilised before intake.

The term improving the nutritional value of an animal feed means improving the availability of the proteins, thereby

improving the growth rate and/or weight gain and/or feed conversion (i.e. the weight of feed required to produce a unit weight of animal).

The protease can be added to the feed in any form, be it as a relatively pure protease, or in admixture with other components intended for addition to animal feed, i.e. in the form of animal feed additives, such as the so-called pre-mixes for animal feed.

Animal feed additives

Apart from the acid-stable protease, the animal feed additives of the invention contain at least one fat-soluble vitamin, and/or at least one water-soluble vitamin, and/or at least one trace mineral, and/or at least one macro mineral.

Further, optional, feed-additive ingredients are colouring agents, aroma compounds, stabilisers, and/or at least one other enzyme selected from amongst phytases EC 3.1.3.8 or 3.1.3.26; xylanases EC 3.2.1.8; galactanases EC 3.2.1.89; and/or beta-glucanases EC 3.2.1.4 (EC refers to Enzyme Classes according to Enzyme Nomenclature 1992 from NC-IUBMB, 1992), see also the World Wide Web (WWW) at <http://www.chem.qmw.ac.uk/iubmb/enzyme/index.html>.

In a particular embodiment these other enzymes are well-defined (as defined and exemplified above for protease preparations, i.a. by reference to Example 8).

Usually fat- and water-soluble vitamins, as well as trace minerals form part of a so-called premix intended for addition to the feed, whereas macro minerals are usually separately added to the feed. Either of these composition types, when enriched with an acid-stable protease according to the invention, is an animal feed additive of the invention.

In a particular embodiment, the animal feed additive of the invention is intended for being included (or prescribed as having to be included) in animal diets or feed at levels of 0.01-10.0%; more particularly 0.05-5.0% or 0.2-1.0% w/w.

Accordingly, the concentration of the individual components of the animal feed additive, and the premix, can be determined by the skilled person in the art.

same component by, respectively, 10-10000; 20-2000; or 100-500 (referring to the above three percentage inclusion intervals).

Guidelines for desired final concentrations, i.e. in-feed-concentrations, of such individual feed and feed additive components are indicated in Table A below.

The following are non-exclusive lists of examples of these components:

Examples of fat-soluble vitamins are vitamin A, vitamin D3, vitamin E, and vitamin K, e.g. vitamin K3.

10 Examples of water-soluble vitamins are vitamin B12, biotin and choline, vitamin B1, vitamin B2, vitamin B6, niacin, folic acid and panthothenate, e.g. Ca-D-panthothenate.

Examples of trace minerals are manganese, zinc, iron, copper, iodine, selenium, and cobalt.

15 Examples of macro minerals are calcium, phosphorus and sodium.

The nutritional requirements of these components - exemplified with poultry and piglets/pigs - are listed in Table A below. Nutritional requirement means that these components should be provided in the diet in the concentrations indicated. These data are compiled from:

NRC, Nutrient requirements in swine, ninth revised edition 1988, subcommittee on swine nutrition, committee on animal nutrition, board of agriculture, national research council. National Academy Press, Washington, D.C. 1988; and

NRC, Nutrient requirements of poultry, ninth revised edition 1994, subcommittee on poultry nutrition, committee on animal nutrition, board of agriculture, national research council. National Academy Press, Washington, D.C. 1994.

30 In the alternative, the animal feed additive of the invention comprises at least one of the individual components specified in Table A. At least one means either of, one or

more of the individual components specified in Table A. The individual component is included in the additive of the invention in such

range indicated in column four, or column five, or column six of Table A.

As explained above, corresponding feed additive concentrations can be found by multiplying the interval limits of these ranges with 10-10000; 20-2000; or 100-500. As an example, considering which premix-content of vitamin A would correspond to the feed-content of 10-10000 IU/kg, this exercise would lead to the following intervals: 100-10⁸ IU; or 200-2x10⁷ IU; or 1000-5x10⁶ IU per kg additive.

Table A

Nutrient requirements - and preferred ranges

Nutrients provided per kg diet	Poultry	Pig-lets/Pigs /Sows	Range 1	Range 2	Range 3
Fat-soluble vitamins					
Vitamin A/[IU]	-5000	1300-4000	10-10000	50-8000	100-6000
Vitamin D ₃ /[IU]	-1100	150-200	2-3000	5-2000	10-1500
Vitamin E/[IU]	-12	11-22	0.02-100	0.2-80	0.5-50
Vitamin K/[mg]	0.5-1.5	-0.5	0.005-10.0	0.05-5.0	0.1-3.0
Water-soluble vitamins					
B ₁₂ /[mg]	-0.003	0.005-0.02	0.0001-1.000	0.0005-0.500	0.001-0.100
Biotin/[mg]	0.100-0.25	0.05-0.08	0.001-10.00	0.005-5.00	0.01-1.00
Choline/[mg]	800-1600	300-600	1-10000	5-5000	10-3000
Trace minerals					
Manganese/[mg]	-60	2.0-4.0	0.1-1000	0.5-500	1.0-100
Zinc/[mg]	40-70	50-100	1-1000	5-500	10-300
Iron/[mg]	50-80	40-100	1-1000	5-500	10-300
Copper/[mg]	6-8	3.0-6.0	0.1-1000	0.5-100	1.0-25
Iodine/[mg]	-0.4	-0.14	0.01-100	0.05-10	0.1-1.0
Selenium/[mg]	-0.2	0.10-0.30	0.005-100	0.01-10.0	0.05-1.0
Macro minerals					
Calcium/[g]	8-40	5-9	0.1-200	0.5-150	1-100
Phosphorus, as available	3-6	1.5-6	0.1-200	0.5-150	1-50

Animal feed compositions

Animal feed compositions - and dietary requirements for various animals

(NRC) publications referred to above, poultry and pig diets can be characterised as indicated in Table B below, columns 2-3. Fish diets can be characterised as indicated in column 4 of Table B. Furthermore such fish diets usually have a crude fat
 5 content of 200-310 g/kg. These fish diet are exemplified with diets for Salmonids and designed on the basis of Aquaculture, principles and practices, ed. T.V.R. Pillay, Blackwell Scientific Publications Ltd. 1990; Fish nutrition, second edition, ed. John E. Halver, Academic Press Inc. 1989.

10 An animal feed composition according to the invention has a crude protein content of 50-800 g/kg, and furthermore comprises at least one protease as claimed herein.

Furthermore, or in the alternative (to the crude protein content indicated above), the animal feed composition of the
 15 invention has a content of metabolisable energy of 10-30 MJ/kg; and/or a content of calcium of 0.1-200 g/kg; and/or a content of available phosphorus of 0.1-200 g/kg; and/or a content of methionine of 0.1-100 g/kg; and/or a content of methionine plus cysteine of 0.1-150 g/kg; and/or a content of
 20 lysine of 0.5-50 g/kg.

In particular embodiments, the content of metabolisable energy, crude protein, calcium, phosphorus, methionine, methionine plus cysteine, and/or lysine is within any one of ranges 2, 3, 4 or 5 in Table B below (R. 2-5).

25 Crude protein is calculated as nitrogen (N) multiplied by a factor 6.25, i.e. Crude protein (g/kg) = N (g/kg) x 6.25 as stated in Animal Nutrition, 4th edition, Chapter 13 (Eds. P. McDonald, R. A. Edwards and J. F. D. Greenhalgh, Longman Scientific and Technical, 1988, ISBN 0-582-40903-9). The ni-
 30 trogen content is determined by the Kjeldahl method (A.O.A.C., 1984, Official Methods of Analysis 14th ed., Association of Official Analytical Chemists, Washington DC).

The author is grateful to the following for their help and support: Dr. J. H. Spelderholt, Centre for poultry research and extension, Wageningen, The Netherlands; Dr. J. H. Spelderholt, Wageningen, The Netherlands; Dr. J. H. Spelderholt, Wageningen, The Netherlands.

The dietary content of calcium, available phosphorus and amino acids in complete animal diets is calculated on the basis of feed tables such as Veevoedertabel 1997, gegevens over chemische samenstelling, verteerbaarheid en voederwaarde van voedermiddelen, Central Veevoederbureau, Runderweg 6, 8219 pk Lelystad. ISBN 90-72839-13-7.

In a particular embodiment, the animal feed composition of the invention contains at least one vegetable protein or protein source as defined above.

10 In still further particular embodiments, the animal feed composition of the invention contains 0-80% maize; and/or 0-80% sorghum; and/or 0-70% wheat; and/or 0-70% Barley; and/or 0-30% oats; and/or 0-40% soybean meal; and/or 0-10% fish meal; and/or 0-20% whey.

15 Animal diets can e.g. be manufactured as mash feed (non-pelleted) or pelleted feed. Typically, the milled feed-stuffs are mixed and sufficient amounts of essential vitamins and minerals are added according to the specifications for the species in question. Enzymes can be added as solid or liquid
20 enzyme formulations. For example, a solid enzyme formulation is typically added before or during the mixing step; and a liquid enzyme preparation is typically added after the pelleting step. The enzyme may also be incorporated in a feed additive or premix. The final enzyme concentration in the diet is
25 within the range of 0.01-200 mg enzyme protein per kg diet, for example in the range of 5-30 mg enzyme protein per kg animal diet.

Examples of animal feed compositions are shown in Example 7.

Table B

Range values for energy, protein and minerals in animal diets

Nutrient	Poul- try	Pig- lets/Pigs /Sows	Fish	R. 1	R. 2	R. 3	R. 4	R. 5
	Min - Max	Min - Max	Min - Max					
Metabo- lisable energy, MJ/kg	12.1 - 13.4	12.9-13.5	14- 25	10- 30	11- 28	11- 26	12- 25	
Crude protein, g/kg	124- 280	120-240	300- 480	50- 800	75- 700	100- 600	110- 500	120- 490
Calcium, g/kg	8-40	5-9	10- 15	0.1- 200	0.5- 150	1- 100	4-50	
Avail- able Phospho- rus, g/kg	2.1- 6.0	1.5-5.5	3-12	0.1- 200	0.5- 150	1- 100	1-50	1-25
Methio- nine, g/kg	3.2 -5.5	-	12- 16	0.1- 100	0.5- 75	1-50	1-30	
Methio- nine plus Cys- teine, g/kg	4-9	2.3-6.8	-	0.1- 150	0.5- 125	1-80		
Lysine, g/kg	2.5- 11	6-14	12- 22	0.5- 50	0.5- 40	1-30		

5

In particular embodiments of the method of the invention for treating vegetable proteins, a further step of adding phytase is also included. And in further particular 10 embodiments, in addition to the combined treatment with phytase and protease, further enzymes may also be added, wherein these enzymes are selected from the group comprising

protease which can be applied in an effective amount, i.e. in an amount adequate for improving

is at present contemplated that the enzyme is administered in one or more of the following amounts (dosage ranges): 0.01-200; or 0.01-100; or 0.05-100; or 0.05-50; or 0.10-10 - all these ranges being in mg protease protein per kg feed (ppm).

5 For determining mg protease protein per kg feed, the protease is purified from the feed composition, and the specific activity of the purified protease is determined using a relevant assay (see under protease activity, substrates, and assays). The protease activity of the feed composition as such
10 is also determined using the same assay, and on the basis of these two determinations, the dosage in mg protease protein per kg feed is calculated.

The same principles apply for determining mg protease protein in feed additives.

15 Of course, if a sample is available of the protease used for preparing the feed additive or the feed, the specific activity is determined from this sample (no need to purify the protease from the feed composition or the additive).

Many vegetables contain anti-nutritional factors such as
20 lectins and trypsin inhibitors. The most important anti-nutritional factors of soybean are the lectin soybean agglutinin (SBA), and the soybean trypsin inhibitor (STI).

Lectins are proteins that bind to specific carbohydrate-containing molecules with considerable specificity, and when
25 ingested they become bound to the intestinal epithelium. This may lead to reduced viability of the epithelial cells and reduced absorption of nutrients.

SBA is a glycosylated, tetrameric lectin with a subunit molecular weight of about 30 kDa and a high affinity for N-
30 acetylgalactosamine.

Trypsin inhibitors affect the intestinal proteolysis reducing protein digestibility and also decrease the

example of a trypsin inhibitor is the Bowman-Birk inhibitor, that has a molecular weight of about 8 kDa, contains 7

are the so-called Kunitz Inhibitors of Factors (e.g. the Soybean Kunitz Trypsin Inhibitor that contains one binding site for trypsin-like proteases and has a molecular weight of about 20 kDa).

5 The proteases for use according to the invention have been shown to hydrolyse anti-nutritional factors like SBA lectin, and the trypsin inhibitors Bowman Birk Inhibitor and The Soybean Kunitz Factor. See the experimental part, Example 5.

10 Thus, the invention also relates to the use of acid-stable proteases for hydrolysing, or reducing the amount of, anti-nutritional factors, e.g. SBA lectin, and trypsin inhibitors, such as the Bowman Birk Inhibitor, and Kunitz Factors, such as the Soybean Kunitz Factor.

15

Example 1

Screening for acid-stable proteases

A large number of proteases were analysed for stability at pH 3, with the objective of identifying proteases that have
20 the necessary stability to pass through the acidic stomach of mono-gastric animals.

The proteases had been purified by conventional chromatographic methods such as ion-exchange chromatography, hydrophobic interaction chromatography and size exclusion
25 chromatography (see e.g. Protein Purification, Principles, High Resolution Methods, and Applications. Editors: Jan-Christer Janson, Lars Rydén, VCH Publishers, 1989).

Protease activity was determined as follows: The protease was incubated with 1.67% Hammarsten casein at 25°C,
30 pH 9.5 for 30 minutes, then TCA (tri-chloro acetic acid) was added to a final concentration of 2% (w/w), the mixture was filtrated to remove the sediment and the filtrate was

measured for absorbance at 280 nm, using a protein standard
krochemische Taschenbuch Teil II, Springer Verlag (1964),

amino groups per minute under standard conditions, i.e. 25°C and pH 9.5.

The proteases were diluted to an activity of 0.6 CPU/l in water, divided in two aliquots and each aliquot was then further diluted to 0.3 CPU/l with 100 mM citrate buffer, pH 3, and 100 mM phosphate buffer, pH 7 respectively. The diluted samples were incubated at 37°C for 1 hour, and 20 µl of the samples were applied to holes in 1% agarose plates containing 1% skim milk. The plates (pH 7.0) were incubated at 37°C over night and clearing zones were measured.

A number of proteases performed well in this test, and the following two have now been characterised: the *Nocardiosis alba* and *Nocardiosis* sp. NRRL 18262 proteases described in example 2. The strain of *Nocardiosis alba* has been deposited according to the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure at DSMZ-Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH, Mascheroder Weg 1b, D-38124 Braunschweig, Germany, as follows.

20

Deposit date	:	22 January 2001
DSM No.	:	<i>Nocardiosis alba</i> DSM 14010

The deposit was made by Novozymes A/S and was later assigned to Hoffmann-La Roche AG

Example 2

Preparation, characterisation and comparative study of *Nocardiosis* proteases

Fermentation

Nocardiosis alba was inoculated from tryptone yeast

medium (tryptone 10 g/l, yeast extract 5 g/l, KH_2PO_4 0.5 g/l, $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ 0.5 g/l, NaCl 0.5 g/l, H_2 -hydrogen phosphate 0.5 g/l, Fluoronic PB 6100 0.2 ml/l in 1000 ml water) into 200 ml of the same medium in a 250 ml flask.

Purification

The culture broth was centrifuged at 10000 x g for 30 minutes in 1 liter beakers. The supernatants were combined and further clarified by a filtration through a Seitz K-250 depth filter plate. The clear filtrate was concentrated by ultrafiltration on a 3kDa cut-off polyether sulfone cassette (Filtron). The concentrated enzyme was transferred to 50mM H₃BO₃, 5mM 3,3'-dimethyl glutaric acid, 1mM CaCl₂, pH 7 (Buffer A) on a G25 Sephadex column (Amersham Pharmacia Biotech), and applied to a Bacitracin agarose column (Upfront Chromatography A/S) equilibrated in Buffer A. After washing the Bacitracin column with Buffer A to remove unbound protein, the protease was eluted from the column using Buffer A supplemented with 25% 2-propanol and 1M sodium chloride. The fractions with protease activity were pooled and transferred to 20mM CH₃COOH/NaOH, 1mM CaCl₂, pH 4.5 (Buffer B) by chromatography on a G25 Sephadex column (Amersham Pharmacia Biotech). The buffer exchanged protease pool was applied to a SOURCE 30S column (Amersham Pharmacia Biotech) equilibrated in Buffer B. After washing the SOURCE 30S column with Buffer B, the protease was eluted with an increasing linear NaCl gradient (0 to 0.25M) in Buffer B. Fractions from the column were tested for protease activity and protease containing fractions were analysed by SDS-PAGE. Pure fractions were pooled and used for further characterisation.

The protease of *Nocardiopsis* sp. NRRL 18262 was prepared using conventional methods, as generally described above for the protease of *Nocardiopsis*.

Characterisation

The protease derived from *Nocardiopsis alba* was found to have a molecular weight of Mr = 21 kDa (SDS-PAGE), and the following partial (N-terminal (MVS)) amino acid sequence was

ADLIDGLAYTMGGEC SVGF AATNAAGQPGFVTAGHCDFVETQVTIGDERGVFEQSV

ADLIDGLAYTMGGEC SVGF AATNAAGQPGFVTAGHCDFVETQVTIGDERGVFEQSV

QSVSYPEGTVTNMTRTTVCAEPGDSGGSYISGTQAQGVTSGGSGNCRTGGTTTFYQEVTPMVN
 SWGVRLRT (SEQ ID NO: 1).

The purpose of this characterisation was to study their
 pH-stability, pH-activity and temperature-activity profiles,
 5 in comparison to Sub.Novo, Sub.Novo(Y217L), and SAVINASE™.

Sub.Novo is subtilisin from *Bacillus amyloliquefaciens*,
 and Sub.Novo(Y217L) is the mutant thereof that is disclosed in
 WO96/05739. Sub.Novo was prepared and purified from a culture
 of the wild-type strain using conventional methods, whereas
 10 the mutant was prepared as described in Examples 1-2, and 15-
 16 of EP 130756.

SAVINASE™ is a subtilisin derived from *Bacillus clausii*
 (previously *Bacillus lentus* NCIB 10309), commercially avail-
 able from Novozymes A/S, Krogshoejvej, DK-2880 Bagsvaerd, Den-
 15 mark. Its preparation is described in US patent No. 3723250.

Example 2A

Determination of SDS-PAGE purity of protease samples

The SDS-PAGE purity of the protease samples was
 20 determined by the following procedure:

40µl protease solution (A_{280} concentration = 0.025) was
 mixed with 10µl 50%(w/v) TCA (trichloroacetic acid) in an Ep-
 ENDORF tube on ice. After half an hour on ice the tube was
 centrifuged (5 minutes, 0°C, 14.000 x g) and the supernatant
 25 was carefully removed. 20µl SDS-PAGE sample buffer (200µl
 Tris-Glycine SDS Sample Buffer (2x) (125mM Tris/HCl, pH 6.8,
 4%(w/v) SDS, 50ppm bromophenol blue, 20%(v/v) Glycerol, LC2676
 from NOVEX™) + 160µl dist. water + 20µl β-mercaptoethanol +
 20µl 3M unbuffered Tris Base (Sigma T-1503) was added to the
 30 precipitate and the tube was boiled for 3 minutes. The tube

was then cooled and the sample was separated on a 15% SDS-PAGE gel.
 the gel, (Laemmli, U.K., (1970) Nature, vol. 227, pp. 680-685).

SDS, distilled water to 1 liter) in both buffer reservoirs at a 150V constant voltage until the bromophenol blue tracking dye had reached the bottom of the gel. After electrophoresis, the gel was rinsed 3 times, 5 minutes each, with 100 ml of distilled water by gentle shaking. The gel was then gently shaken with Gelcode[®] Blue Stain Reagent (colloidal Coomassie G-250 product from PIERCE, PIERCE cat. No. 24592) for one hour and washed by gentle shaking for 8 to 16 hours with distilled water with several changes of distilled water. Finally, the gel was dried between 2 pieces of cellophane. Dried gels were scanned with a Arcus II scanner from AGFA equipped with Fotolook 95 v2.08 software and imported to the image evaluation software CREAM[™] for Windows (catalogue nos. 990001 and 990005, Kem-En-Tec, Denmark) by the *File/Acquire* command with the following settings (of Fotolook 95 v2.08): Original=Reflective, Mode=Color RGB, Scan resolution=240 ppi, Output resolution=1201pi, Scale factor=100%, Range=Histogram with Global selection and Min=0 and Max=215, ToneCurve=None, Sharpness=None, Descreen=None and Flavor=None, thereby producing an *.img picture file of the SDS-PAGE gel, which was used for evaluation in CREAM[™]. The *.img picture file was evaluated with the menu command *Analysis/1-D*. Two scan lines were placed on the *.img picture file with the *Lane Place Tool*: A Sample scan line and a Background scan line. The Sample scan line was placed in the middle of a sample lane (with the protease in question) from just below the application slot to just above the position of the Bromphenol blue tracking dye. The Background scan line was placed parallel to the Sample scan line, but at a position in the pictured SDS-PAGE gel where no sample was applied, start and endpoints for the Background scan line were perpendicular to the start and endpoints of the Sample

scanned with the *1-D Scan* menu command with *Medium* sensitivity. Using the *1-D Filter* menu command the background scan line

command was selected and the Area % of the protease peak, as calculated by the CREAM™ software, was used as the SDS-PAGE purity of the proteases.

All the protease samples had an SDS-PAGE purity of above
5 95%.

Example 2B

pH-activity assay

Suc-AAPF-pNA (Sigma® S-7388) was used for obtaining pH-
10 activity profiles.

Assay buffer: 100mM succinic acid (Merck 1.00682), 100mM HEPES (Sigma H-3375), 100mM CHES (Sigma C-2885), 100mM CABS (Sigma C-5580), 1mM CaCl₂, 150mM KCl, 0.01% Triton® X-100, adjusted to pH-values 3.0, 4.0, 5.0, 6.0, 7.0, 8.0, 9.0, 10.0,
15 or 11.0 with HCl or NaOH.

Assay temperature: 25°C.

A 300µl protease sample (diluted in 0.01% Triton® X-100) was mixed with 1.5 ml of the assay buffer at the respective pH value, bringing the pH of the mixture to the pH of the assay
20 buffer. The reaction was started by adding 1.5ml pNA substrate (50mg dissolved in 1.0ml DMSO and further diluted 45x with 0.01% Triton® X-100) and, after mixing, the increase in A₄₀₅ was monitored by a spectrophotometer as a measurement of the protease activity at the pH in question. The assay was
25 repeated with the assay buffer at the other pH values, and the activity measurements were plotted as relative activity against pH. The relative activities were normalized with the highest activity (pH-optimum), i.e. setting activity at pH-optimum to 1, or to 100%. The protease samples were diluted to
30 ensure that all activity measurements fell within the linear part of the dose-response curve for the assay.

Example 2CpH-stability assay

Suc-AAPF-pNA (Sigma® S-7388) was used for obtaining pH-stability profiles.

5 Assay buffer: 100mM succinic acid, 100mM HEPES, 100mM CHES, 100mM CABS, 1mM CaCl_2 , 150mM KCl, 0.01% Triton® X-100 adjusted to pH-values 2.0, 2.5, 3.0, 3.5, 4.0, 4.5, 5.0, 6.0, 7.0, 8.0, 9.0, 10.0 or 11.0 with HCl or NaOH.

Each protease sample (in 1mM succinic acid, 2mM CaCl_2 ,
10 100mM NaCl, pH 6.0 and with an A_{280} absorption > 10) was diluted in the assay buffer at each pH value tested to $A_{280} = 1.0$. The diluted protease samples were incubated for 2 hours at 37°C. After incubation, protease samples were diluted in 100mM succinic acid, 100mM HEPES, 100mM CHES, 100mM CABS, 1mM CaCl_2 ,
15 150mM KCl, 0.01% Triton® X-100, pH 9.0, bringing the pH of all samples to pH 9.0.

In the following activity measurement, the temperature was 25°C.

300 μ l diluted protease sample was mixed with 1.5ml of
20 the pH 9.0 assay buffer and the activity reaction was started by adding 1.5ml pNA substrate (50mg dissolved in 1.0ml DMSO and further diluted 45x with 0.01% Triton® X-100) and, after mixing, the increase in A_{405} was monitored by a spectrophotometer as a measurement of the (residual) protease activity.
25 The 37°C incubation was performed at the different pH-values and the activity measurements were plotted as residual activities against pH. The residual activities were normalized with the activity of a parallel incubation (control), where the protease was diluted to $A_{280} = 1.0$ in the assay buffer at pH
30 9.0 and incubated for 2 hours at 5°C before activity measurement as the other incubations. The protease samples

Example 2DTemperature-activity assay

Protazyme AK tablets were used for obtaining temperature profiles. Protazyme AK tablets are azurine dyed crosslinked casein prepared as tablets by Megazyme.

Assay buffer: 100mM succinic acid, 100mM HEPES, 100mM CHES, 100mM CABS, 1mM CaCl_2 , 150mM KCl, 0.01% Triton[®] X-100 adjusted to pH 9.0 with NaOH.

A Protazyme AK tablet was suspended in 2.0ml 0.01% Triton[®] X-100 by gentle stirring. 500 μ l of this suspension and 500 μ l assay buffer were mixed in an Eppendorf tube and placed on ice. 20 μ l protease sample (diluted in 0.01% Triton X-100) was added. The assay was initiated by transferring the Eppendorf tube to an Eppendorf thermomixer, which was set to the assay temperature. The tube was incubated for 15 minutes on the Eppendorf thermomixer at its highest shaking rate. By transferring the tube back to the ice bath, the assay incubation was stopped. The tube was centrifuged in an ice-cold centrifuge for a few minutes and the A_{650} of the supernatant was read by a spectrophotometer. A buffer blind was included in the assay (instead of enzyme). $A_{650}(\text{protease}) - A_{650}(\text{blind})$ was a measurement of protease activity. The assay was performed at different temperatures and the activity measurements were plotted as relative activities against incubation temperature. The relative activities were normalized with the highest activity (temperature optimum). The protease samples were diluted to ensure that all activity measurements fell within the near linear part of the dose-response curve for the assay.

An overview of the activity optima (pH- and temperature activity) is seen in Table 1. pH-stability, pH-activity and temperature-activity profiles are seen in figures 1-3, and a

Table 1

pH- and temperature optima of various proteases

Protease	pH-optimum (pNA-substrate)	Temperature-optimum at pH 9.0 (Protazyme AK)
Nocardiosis sp. NRRL 18262	10	70°C
Nocardiosis alba	11	70°C
Sub.Novo ¹	10	70°C
Sub.Novo (Y217L) ²	9	70°C
SAVINASE ^{TM3}	9	70°C

5 Table 2

pH-stability of various proteases, between pH 2.0 and 5.0

Protease	pH 2.0	pH 2.5	pH 3.0	pH 3.5	pH 4.0	pH 4.5	pH 5.0
Nocardiosis sp. NRRL 18262	0.779	1.000	1.029	0.983	0.991	1.019	1.004]
Nocardiosis alba	0.929	0.993	1.009	1.005	0.969	1.037	0.992
Sub.Novo	0.007	0.003	0.000	0.000	0.024	0.784	0.942
Sub.Novo (Y217L)	0.000	0.000	0.002	0.003	0.350	0.951	0.996
Savinase	0.001	0.001	0.001	0.003	0.338	0.929	0.992

Example 2E

10 Absorption purity of purified protease samples

Determination of A_{280}/A_{260} ratio

The A_{280}/A_{260} ratio of purified protease samples was determined as follows.

A_{260} means the absorption of a protease sample at 260 nm in a 1cm path length cuvette relative to a buffer blank. A_{280} means the absorption of the same protease sample at 280 nm in a 1cm path length cuvette relative to a buffer blank.

Samples of the purified proteases from Example 2 were

1. Nocardiosis sp. NRRL 18262, 2. Nocardiosis alba, 3. Sub.Novo, 4. Sub.Novo (Y217L), 5. Savinase

Example 3Ability of protease derived from Nocardiosis sp. NRRL 18262 to degrade insoluble parts of Soy Bean Meal (SBM)

5 The protease from Nocardiosis sp. NRRL 18262 was tested for its ability to make the insoluble/indigestible parts of SBM accessible to digestive enzymes and/or added exogenous enzymes.

 Its performance was compared to two aspartate proteases, 10 Protease I and Protease II, prepared as described in WO 95/02044. This document also discloses their use in feed. Protease I is an Aspergillopepsin II type of protease, and Protease II an Aspergillopepsin I type of protease (both aspartate proteases, i.e. non-subtilisin proteases) from 15 Aspergillus aculeatus (reference being made to Handbook of Proteolytic Enzymes referred to above).

 The test substrate, the so-called soy remnant, was produced in a process which mimics the digestive tract of mono-gastric animals, including a pepsin treatment at pH 2, 20 and a pancreatin treatment at pH 7.

 In the pancreatin treatment step a range of commercial enzymes was added in high dosages in order to degrade the SBM components that are accessible to existing commercial enzymes.

 The following enzymes, all commercially available from 25 Novozymes A/S, Denmark, were added: ALCALASE™ 2.4L, NEUTRASE™ 0.5L, FLAVOURZYME™ 1000L, ENERGEX™ L, BIOFEED™ Plus L, PHYTASE NOVO™ L. The SBM used was a standard 48% protein SBM for feed, which had been pelletised.

 After the treatment only 5% of the total protein was 30 left in the resulting soy remnant.

FITC labelling protocol

 The SBM was suspended in water and stirred for 1 hour at 40 °C. The suspension was cooled to room temperature and treated with fluorescein-5-isothiocyanate

(FITC) over night in the dark. Non-coupled probe was removed by ultrafiltration (10.000 Mw cut-off).

FITC-assay

The FITC-labelled soy remnant was used for testing the ability of the proteases to degrade the soy remnant using the following assay: 0.4 ml protease sample (with $A_{280} = 0.1$) was mixed with 0.4 ml FITC-soy remnant (suspension of 10 mg/ml in 0.2M sodium-phosphate buffer pH 6.5) at 37°C, and the relative fluorescence units (RFU 485/535nm; excitation/monitoring wavelength) measured after 0 hours, and after 22 hours incubation. Before determination of the RFU, samples were centrifuged for 1 min at 20.000 x G and 250 micro-liter supernatant was transferred to a black micro-titer tray. Measurements were performed using a VICTOR 1420 Multilabel counter (*In vitro*, Denmark). RFU is generally described by Iain D. Johnson in: Introduction to Fluorescence Techniques, Handbook of Fluorescent Probes and Research Chemicals, Molecular Probes, Richard P. Haugland, 6th edition, 1996 (ISBN 0-9652240-0-7).

A blind sample was prepared by adding 0.4 ml buffer instead of enzyme sample.

$RFU_{sample} = \Delta RFU_{sample} - \Delta RFU_{blind}$, where $\Delta RFU = RFU(22 \text{ hours}) - RFU(0 \text{ hours})$

The resulting FITC values (RFU_{sample} values) are shown in Table 3 below. The FITC values are generally with an error margin of +/- 20.000. Contrary to Protease I and Protease II, the protease derived from *Nocardiosis* sp. NRRL 18262 degraded the soy remnant to a significant extent.

Table 3

Ability of proteases to degrade soy remnant

Protease	FITC (+/-20000)
<i>Nocardiosis</i> sp. NRRL 18262	92900

Example 4In vitro testing of the protease derived from Nocardiosis sp. NRRL 18262

The protease derived from Nocardiosis sp, NRRL 18262 was tested, together with a protease derived from Bacillus sp. NCIMB 40484 ("PD498", prepared as described in Example 1 of WO93/24623), and together with FLAVOURZYME™, a protease-containing enzyme preparation from Aspergillus oryzae (commercially available from Novozymes A/S, Bagsvaerd, Denmark), for its ability to solubilise maize-SBM (maize-Soy Bean Meal) proteins in an in vitro digestion system (simulating digestion in monogastric animals). For the blank treatments, maize-SBM was incubated in the absence of exogenous proteases.

15 Outline of in vitro model

Components added to flask	Time course (min)
10g maize-SBM (60:40) + HCl/pepsin (3000U/g diet) + protease (0.1mg enzyme protein/g diet or 3.3 mg FLAVOURZYME™ /g diet), T=40°C, pH=3.0	t=0
NaOH, T=40°C, pH 6	t=60
NaHCO ₃ /pancreatin (8.0mg/g diet), T=40°C, pH 6-7	t=80
Stop incubation and take samples, T=0°C	t=320

Substrates

10 g maize-SBM diet with a ratio maize-SBM of 6:4 (w/w) was used. The protein content was 43% (w/w) in SBM and 8.2% (w/w) in maize meal. The total amount of protein in 10 g maize-SBM diet was 2.21 g.

Digestive enzymes

Pepsin (Sigma P-7000; 539 U/mg, solid), pancreatin (Sigma P-7545; 8xU.S.P. (US Pharmacopeia))

The ratio of the A₁ and A₂ and the amino acid composition and acid composition using the principles outlined in S.C.Gill &

Experimental procedure for *in vitro* model

1. 10 g of substrate is weighed into a 100 ml flask.
2. At time 0 min, 46 ml HCl (0.1 M) containing pepsin (3000 U/g diet) and 1 ml of protease (0.1 mg enzyme protein/g diet, except for FLAVOURZYME™: 3.3 mg/g diet) are added to the flask while mixing. The flask is incubated at 40°C.
3. At time 20-25 min, pH is measured.
4. At time 45 min, 16 ml of H₂O is added.
5. At time 60 min, 7 ml of NaOH (0.4 M) is added.
- 10 6. At time 80 min, 5 ml of NaHCO₃ (1 M) containing pancreatin (8.0 mg/g diet) is added.
7. At time 90 min, pH is measured.
8. At time 300 min, pH is measured.
9. At time 320 min, aliquots of 30 ml are removed and
15 centrifuged (10000 x g, 10 min, 0°C).
10. Total soluble protein in supernatants is determined.

Protein determination

Supernatants are analysed for protein content using the Kjeldahl method (determination of % nitrogen; A.O.A.C. (1984) Official Methods of Analysis 14th ed. Association of Official Analytical Chemists, Washington DC).

Calculations

For all samples, *in vitro* protein solubility was calculated using the equations below.

25 Amount of protein in diet sample: 22.1% of 10 g = 2.21 g

If all the protein were solubilised in the 75 ml of liquid, the protein concentration in the supernatant would be: 2.21 g/75 ml ≈ 2.95%.

Note that the supernatants also include the digestive
30 and exogenous enzymes. In order to determine the solubility, the protein contribution from the digestive and exogenous

$((X \text{ mg pancreatin/g diet} \times 10 \text{ g diet} \times 0.69 \times 100\%)/(1000 \text{ mg/g} \times 75 \text{ g})) + ((Y \text{ U pepsin/g diet} \times 10 \text{ g diet} \times 0.57 \times 100\%)/(539 \text{ U/mg} \times 1000 \text{ mg/g} \times 75 \text{ g})),$

where 0.69 and 0.57 refer to the protein contents in the pancreatin and pepsin preparations used (i.e. 69% of the pancreatin, and 57% of the pepsin is protein as determined by the Kjeldahl method referred to above).

% protein from exogenous enzymes (Z mg EP/g diet) = $(Z \text{ mg EP/g diet} \times 10 \text{ g diet} \times 100\%)/(1000 \text{ mg/g} \times 75 \text{ g})$

% protein corrected in supernatant = % protein in supernatant as analysed - (% protein from digestive enzymes + % protein from exogenous enzymes)

Protein solubilisation (%) = $(\% \text{ protein corrected in supernatant} \times 100\%)/2.95 \%$

The results below show that the protease derived from *Nocardiosis* sp. NRRL 18262 has a significantly better effect on protein solubilisation as compared to the blank, and as compared to the *Bacillus* sp. NCIMB 40484 protease.

Enzyme	Solubilised P (% of total)	SD	n
Blind (no exogenous enzymes)	73.8 ^c	0.87	10
+ the protease derived from <i>Nocardiosis</i> sp. NRRL 18262	77.5 ^a	0.50	10
+ the protease derived from <i>Bacillus</i> sp. NCIMB 40484	75.6 ^b	1.52	5
+ FLAVOURZYME™	74.1 ^c	0.23	4

a, b, c Values within a column not sharing a common superscript letter are significantly different, $P < 0.05$. SD is standard deviation; n is the number of observations.

The ability of the proteases from *Nocardiosis* sp. NRRL 18262 and *Bacillus* sp. NCIMB 40484 to hydrolyse soybean

agglutinin (SBA) and the soy Bowman-Birk and Kunitz trypsin inhibitors was tested.

Pure SBA (Fluka 61763), Bowman-Birk Inhibitor (Sigma T-9777) or Kunitz Inhibitor (Trypsin Inhibitor from soybean, 5 Boehringer Mannheim 109886) was incubated with the protease for 2 hours, 37°C, at pH 6.5 (protease: anti-nutritional factor = 1:10, based on A_{280}). Incubation buffer: 50 mM dimethyl glutaric acid, 150 mM NaCl, 1 mM CaCl_2 , 0.01% Triton X-100, pH 6.5.

10 The ability of the proteases to degrade SBA and the protease inhibitors was estimated from the disappearance of the native SBA or trypsin inhibitor bands and appearance of low molecular weight degradation products on SDS-PAGE gels. Gels were stained with Coomassie blue and band intensity 15 determined by scanning.

The results, as % of anti-nutritional factor degraded, are shown in Table 4 below.

It is contemplated that the ability to degrade the anti-nutritional factors in soy can also be estimated by applying 20 the Western technique with antibodies against SBA, Bowman-Birk Inhibitor or Kunitz Inhibitor after incubation of soybean meal with the candidate proteases (see WO98/56260).

Table 4

Protease derived from	SBA	Bowman-Birk Inhibitor	Kunitz Inhibitor
Nocardiosis sp. NRRL 18262	75	25	100
Bacillus sp. NCIMB 40484	21	41	100

Example 6Effects of acid-stable Nocardiosis proteases on the growth performance of broiler chickens

The trial is carried out in accordance with the official French instructions for experiments with live animals. Day-old broiler chickens ('Ross PM3'), separated by sex, are supplied by a commercial hatchery.

The chickens are housed in wire-floored battery cages, which are kept in an environmentally controlled room. Feed and tap water is provided *ad libitum*.

On day 8, the chickens are divided by weight into groups of 6 birds, which are allocated to either the control treatment, receiving the experimental diet without enzymes, or to the enzyme treatment, receiving the experimental diet supplemented with 100 mg enzyme protein of the protease per kg feed.

Each treatment is replicated with 12 groups, 6 groups of each sex. The groups are weighed on days 8 and 29. The feed consumption of the intermediate period is determined and body weight gain and feed conversion ratio are calculated.

The experimental diet is based on maize starch and soybean meal (44 % crude protein) as main ingredients (Table 5). The feed is pelleted (die configuration: 3 x 20 mm) at about 70°C. An appropriate amount of the protease is diluted in a fixed quantity of water and sprayed onto the pelleted feed. For the control treatment, adequate amounts of water are used to handle the treatments in the same way.

For the statistical evaluation, a two factorial analysis of variance (factors: treatment and sex) is carried out, using the GLM procedure of the SAS package (SAS Institute Inc., 1985). When significant treatments effects ($p < 0.05$) are in-

teracted with sex, a two factorial analysis of variance is carried out. In addition, a two factorial analysis of variance is carried out, taking into consideration the main

References

EEC (1986): Directive de la Commission du 9 avril 1986 fixant la méthode de calcul de la valeur énergétique des aliments composés destinés à la volaille. Journal Officiel des Communautés Européennes, L130, 53 - 54

SAS Institute Inc. (1985): SAS® User's Guide, Version 5 Edition. Cary NC

10

Table 5

Composition of the experimental diet

<u>Ingredients (%) :</u>	
Maize starch	45.80
Soybean meal 44 ¹	44.40
Tallow	3.20
Soybean oil	1.00
DL-Methionine	0.18
MCP	0.76
Salt	0.05
Binder	1.00
Vitamin and mineral premix	3.55
Avatec® 15% CC ²	0.06
<u>Analysed content:</u>	
Crude protein (%)	19.3
ME, N-corrected (MJ/kg) ³	12.2
Crude fat (%)	5.3

- 15 ¹ analysed content: 90.6% dry matter, 45.3% crude protein, 2.0% crude fat, 4.9% crude fibre
corresponded to 90 mg lasalocid-Na / kg feed as anticoccidial
² supplemented with 100 mg of avatec 15% CC / kg feed

16 ³ calculated value

Maize starch: 45.80%, Soybean meal: 44.40%, Tallow: 3.20%, Soybean oil: 1.00%, DL-Methionine: 0.18%, MCP: 0.76%, Salt: 0.05%, Binder: 1.00%, Vitamin and mineral premix: 3.55%, Avatec® 15% CC: 0.06%

Tallow: Fondoirs Gachot SA, F-67100 Strasbourg, France
 Soybean oil: Ewoco Sarl, F-68970 Guemar, France
 DL-Methionine: Produit Roche SA, F-92521 Neuilly-sur-Seine,
 France

5 MCP: Brenntag Lorraine, F-54200 Toul, France

Salt: Minoterie Moderne, F-68560 Hirsingue, France

Binder: Minoterie Moderne, F-68560 Hirsingue, France

Premix (AM vol chair NS 4231): Agrobases, F-01007 Bourg-en-
 Bresse, France

10 Avatec: Produit Roche SA, F-92521 Neuilly-sur-Seine, France

Example 7

Premix and diets for turkey and salmonids supplemented with
 acid-stable Nocardiosis proteases.

15 A premix of the following composition is prepared (con-
 tent per kilo):

	5000000	IE	Vitamin A
	1000000	IE	Vitamin D3
	13333	mg	Vitamin E
20	1000	mg	Vitamin K3
	750	mg	Vitamin B1
	2500	mg	Vitamin B2
	1500	mg	Vitamin B6
	7666	mg	Vitamin B12
25	12333	mg	Niacin
	33333	mg	Biotin
	300	mg	Folic Acid
	3000	mg	Ca-D-Panthothenate
	1666	mg	Cu
30	16666	mg	Fe
	16666	mg	Zn

To this premix the protease from *Nocardiopsis* sp. NRRL 18262 is added (prepared as described in Example 2), in an amount corresponding to 10 g protease enzyme protein/kg.

Pelleted turkey starter and grower diets with a composition as shown in the below table (on the basis of Leeson and Summers, 1997 but recalculated without meat meal by using the AGROSOFT®, optimisation program) and with 100 mg protease enzyme protein per kg are prepared as follows:

Milled maize, Soybean meal, Fish-meal and Vegetable fat are mixed in a cascade mixer. Limestone, calcium phosphate and salt are added, together with the above premix in an amount of 10 g/kg diet, followed by mixing. The resulting mixture is pelleted (steam conditioning followed by the pelleting step).

Ingredient	Starter diet, g/kg	Grower, g/kg	Finisher
Maize	454.4	612.5	781.0
Soybean meal	391	279	61.7
Fish meal	70	29.9	70
Vegetable fat	21	21	46
Limestone	19	16.9	9
Calcium phosphate	30	25.9	16.8
Salt (NaCl)	2	2	2
Vitamin and mineral premix	10	10	10
Lysine	1.3	1.49	
Methionine	1.3	1.3	3.6
Calculated nutrients			
Crude protein g/kg	279	213	152
Metabolisable en-	12.3	12.7	14.1

lysine, g/kg			
lysine, g/kg	17.6	12.8	7.5

Two diets for Salmonids are also prepared, as generally outlined above. The actual compositions are indicated in the Table below (compiled from Refstie et al (1998), Aquaculture, vol. 162, p.301-302). The estimated nutrient content is recalculated by using the Agrosoft® feed optimisation program.

The protease derived from *Nocardiosis alba*, prepared as described in Example 2, is added to the diets in an amount corresponding to 100 mg protease enzyme protein per kg.

Ingredient	Conventional diet with fish meal	Alternative diet with soybean meal
Wheat	245.3	75.2
Fish meal	505.0	310.0
Soybean meal	-	339.0
Fish oil	185.0	200.0
DL-Methionine	13.9	23.0
Mono-Calcium phosphate	-	2.0
Vitamin and Mineral premix + pellet binder and astaxanthin	50.8	50.8
Calculated nutrients (fresh weight basis)		
Crude protein g/kg	401	415
Crude fat g/kg	232	247
Metabolisable energy MJ/kg	16.9	16.5
Calcium, g/kg	13.9	9.8
Phosphorus, g/kg	10.8	9.0
Lysine, g/kg	27.7	26.7
Methionine, g/kg	24.4	31.6

size-exclusion column. Size-exclusion chromatography, also known as gel filtration chromatography, is based on a porous gel matrix (packed in a column) with a distribution of pore sizes comparable in size to the protein molecules to be separated. Relatively small protein molecules can diffuse into the gel from the surrounding solution, whereas larger molecules will be prevented by their size from diffusing into the gel to the same degree. As a result, protein molecules are separated according to their size with larger molecules eluting from the column before smaller ones.

Protein concentration assay.

The protein concentration in protease-containing enzyme products is determined with a BCA protein assay kit from PIERCE (identical to PIERCE cat. No.23225). The sodium salt of Bicinchoninic acid (BCA) is a stable, water-soluble compound capable of forming an intense purple complex with cuprous ions (Cu^{1+}) in an alkaline environment. The BCA reagent forms the basis of the BCA protein assay kit capable of monitoring cuprous ions produced in the reaction of protein with alkaline Cu^{2+} (Biuret reaction). The colour produced from this reaction is stable and increases in a proportional fashion with increasing protein concentrations (Smith, P.K., et al. (1985), Analytical Biochemistry, vol. 150, pp. 76-85). The BCA working solution is made by mixing 50 parts of reagent A with 1 part reagent B (Reagent A is PIERCE cat. No. 23223, contains BCA and tartrate in an alkaline carbonate buffer; reagent B is PIERCE cat. No. 23224, contains 4% $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$). 300 μl sample is mixed with 3.0ml BCA working solution. After 30 minutes at 37°C, the sample is cooled to room temperature and A_{490} is read as a measure of the protein concentration in the sample. Dilutions of Bovine serum albumin (PIERCE cat. No. 23209) are included in the assay as a standard.

The protein is first dissolved in 100 μl of 100mM H_2BO_3 , 10mM 3,3' dimethylglutaric acid, 2mM CaCl_2 , pH 6.

a 0.45 μ filter to give a clear solution. The solution is from this point treated as a liquid protease-containing enzyme product.

If the protease-containing enzyme product is a liquid, the product is first dialysed in a 6-8000 Da cut-off SpectraPor dialysis tube (cat.no. 132 670 from Spectrum Medical Industries) against 100 volumes of Buffer A + 150mM NaCl (Buffer B) for at least 5 hours at 5°C, to remove formulation chemicals that could give liquid protease-containing enzyme products a high viscosity, which is detrimental to the size-exclusion chromatography.

The dialysed protease-containing enzyme product is filtered through a 0.45 μ filter if a precipitate was formed during the dialysis. The protein concentration in the dialysed enzyme product is determined with the above described protein concentration assay and the enzyme product is diluted with Buffer B, to give a sample ready for size-exclusion chromatography with a protein concentration of 5 mg/ml. If the enzyme product has a lower than 5 mg/ml protein concentration after dialysis, it is used as is.

Size-exclusion chromatography.

A 300ml HiLoad26/60 Superdex75pg column (Amersham Pharmacia Biotech) is equilibrated in Buffer B (Flow: 1ml/min). 1.0ml of the protease-containing enzyme sample is applied to the column and the column is eluted with Buffer B (Flow: 1ml/min). 2.0ml fractions are collected from the outlet of the column, until all of the applied sample have eluted from the column. The collected fractions are analysed for protein content (see above Protein concentration assay) and for protease activity by appropriate assays. An example of an appropriate assay is the Suc-AAPF-pNA assay (see Example 2B).

described to measure as many proteases in the fractionated sample as possible. The conditions of the assays referred to

conditions are mentioned above in the section dealing with measurement of protease activity. A protein peak with activity in one or more of the protease assays is defined as a protease peak. The purity of a protease peak is calculated as the
5 protein amount in the peak divided with the total protein amount in all identified protease peaks.

The purity of a protease-containing enzyme product is calculated as the amount of protein in the acid-stable protease peak divided with the protein amount in all
10 identified protease peaks using the above procedure.